arXiv:0802.1228v3 [math.NT] 12 May 2009

A generalization of the duality for multiple harmonic sums

Gaku Kawashima

Abstract

The duality is a fundamental property of the finite multiple harmonic sums (MHS). In this paper, we prove a duality result for certain generalizations of MHS which appear naturally as the differences of MHS. We also prove a formula for the differences of these generalized MHS.

Keywords: multiple harmonic sum

1 Introduction

In recent years, multiple harmonic sums (MHS for short) have been studied by theoretical physicists and the duality for MHS was discovered [5, Section 8B]. The same formula has appeared in [2, 6] in the study of arithmetical properties of MHS. It is also used in [3] in order to investigate algebraic relations among multiple zeta values.

Here, we explain the duality for MHS. We call an ordered set of positive integers a multi-index. If we introduce the following nested sums

$$s_{\boldsymbol{\mu}}(n) = \sum_{n=n_1 \geq \cdots \geq n_p \geq 0} \frac{1}{(n_1+1)^{\mu_1} \cdots (n_p+1)^{\mu_p}}, \quad 0 \leq n \in \mathbb{Z}$$

for a multi-index $\boldsymbol{\mu} = (\mu_1, \ldots, \mu_p)$, the duality for MHS is expressed as follows:

$$\sum_{k=0}^{n} (-1)^k \binom{n}{k} s_{\boldsymbol{\mu}}(k) = s_{\boldsymbol{\mu}^*}(n), \quad 0 \leq n \in \mathbb{Z}, \tag{1}$$

where $\boldsymbol{\mu}^*$ is some multi-index determined by $\boldsymbol{\mu}$. For example, we have

$$(1,2,3)^* = (2,2,1,1), \quad (2,2,2)^* = (1,2,2,1) \quad \text{and} \quad (4,1,1)^* = (1,1,1,3)$$

by the diagrams



The lower arrows are in the complementary slots to the upper arrows (see [3, Section 2] for the precise definition of the correspondence $\boldsymbol{\mu} \mapsto \boldsymbol{\mu}^*$). A q-analogue of the identity (1) is also known [1].

In [4], the finite nested sums

$$c_{x_1,\dots,x_p}(n) = \sum_{n=n_1\geq\cdots\geq n_p\geq 0} \frac{x_1^{n_1-n_2}\cdots x_{p-1}^{n_{p-1}-n_p} x_p^{n_p}}{(n_1+1)\cdots(n_{p-1}+1)}, \quad 0 \leq n \in \mathbb{Z}$$

with complex parameters $x_1, \dots, x_p$ were studied in order to derive relations among multiple L-values and a generalization of (1)

$$\sum_{k=0}^{n} (-1)^k \binom{n}{k} c_{x_1,\dots,x_p}(k) = c_{1-x_1,\dots,1-x_p}(n), \quad 0 \leq n \in \mathbb{Z} \tag{2}$$

was proved. To obtain the claim (1) from (2), we have only to note that

$$c_{\underbrace{0,\dots,0,1}_{\mu_1},\dots,\underbrace{0,\dots,0,1}_{\mu_p},0}(n) \quad \text{and} \quad c_{\underbrace{0,\dots,0,1}_{\mu_1},\dots,\underbrace{0,\dots,0,1}_{\mu_{p-1}},\underbrace{0,\dots,0,1}_{\mu_p}}(n)$$

are both equal to $s_{\boldsymbol{\mu}}(n)$. The duality (2) is a consequence of the difference formula

$$(\Delta^k c_{x_1,\dots,x_p})(n) = c_{x_1,\dots,x_p;1-x_1,\dots,1-x_p}(n,k), \quad 0 \leq n,\ k \in \mathbb{Z}.$$

The right-hand side is given by the following nested sums

$$c_{x_1,\dots,x_p;y_1,\dots,y_p}(n,k) = \sum_{\substack{n=n_1\geq\cdots\geq n_p\geq 0\\ k=k_1\geq\cdots\geq k_p\geq 0}} P \\ \times \frac{(x_1^{n_1-n_2}\cdots x_{p-1}^{n_{p-1}-n_p} x_p^{n_p})(y_1^{k_1-k_2}\cdots y_{p-1}^{k_{p-1}-k_p} y_p^{k_p})}{(n_1+k_1+1)\cdots(n_{p-1}+k_{p-1}+1)}, \quad 0 \leq n,\ k \in \mathbb{Z}$$

with complex parameters x_i and y_i for $1 \leq i \leq p$, where

$$P = \binom{n+k}{n}^{-1} \binom{n_1-n_2+k_1-k_2}{n_1-n_2} \cdots \binom{n_{p-1}-n_p+k_{p-1}-k_p}{n_{p-1}-n_p} \binom{n_p+k_p}{n_p}.$$

The symbol Δ denotes the difference operator and its definition is given in Section 2.

In the present paper, we ask whether the duality exists for these more complicated nested sums which appear naturally as the differences of $c_{x_1,\dots,x_p}(n)$. We consider more general nested sums

$$c^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}(n_1,\dots,n_r) = \sum_{\substack{n_1=n_{11}\geq\cdots\geq n_{1p}\geq 0\\ \cdots \\ n_r=n_{r1}\geq\cdots\geq n_{rp}\geq 0}} Q \\ \times \frac{(x_{11}^{n_{11}-n_{12}}\cdots x_{1p-1}^{n_{1p-1}-n_{1p}} x_{1p}^{n_{1p}})\cdots(x_{r1}^{n_{r1}-n_{r2}}\cdots x_{rp-1}^{n_{rp-1}-n_{rp}} x_{rp}^{n_{rp}})}{(n_{11}+\cdots+n_{r1}+t_1)\cdots(n_{1p-1}+\cdots+n_{rp-1}+t_{p-1})}, \\ 0 \leq n_1,\ \dots,\ n_r \in \mathbb{Z}$$

with parameters $t_1, \ldots, t_{p-1} \in \mathbb{C} \setminus \{0, -1, -2, \ldots\}$ and $\boldsymbol{x}_i = (x_{i1}, \ldots, x_{ip}) \in \mathbb{C}^p$ for $1 \leq i \leq r$. We need to find an appropriate Q so that the duality holds. In addition, it is necessary for Q to become P on setting $r = 2$ and $t_1 = \cdots = t_{p-1} = 1$. The following

$$Q = \binom{n_1}{n_{11} - n_{12}, \ldots, n_{1p-1} - n_{1p}, n_{1p}} \cdots \binom{n_r}{n_{r1} - n_{r2}, \ldots, n_{rp-1} - n_{rp}, n_{rp}}$$
$$\times \binom{n_{11} + \cdots + n_{r1} + t_1 - 1}{n_{11} - n_{12} + \cdots + n_{r1} - n_{r2}}^{-1} \cdots \binom{n_{1p-1} + \cdots + n_{rp-1} + t_{p-1} - 1}{n_{1p-1} - n_{1p} + \cdots + n_{rp-1} - n_{rp}}^{-1},$$

which consists of multinomial coefficients and generalized binomial coefficients, satisfies such requirements. In fact, for the above Q we have

$$\sum_{k_1=0}^{n_1} \cdots \sum_{k_r=0}^{n_r} (-1)^{k_1+\cdots+k_r} \binom{n_1}{k_1} \cdots \binom{n_r}{k_r} c_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}^{t_1,\ldots,t_{p-1}}(k_1, \ldots, k_r)$$
$$= c_{\boldsymbol{1}-\boldsymbol{x}_1;\cdots;\boldsymbol{1}-\boldsymbol{x}_r}^{t_1,\ldots,t_{p-1}}(n_1, \ldots, n_r), \quad 0 \leq n_1, \ldots, n_r \in \mathbb{Z},$$

where $\boldsymbol{1} - \boldsymbol{x} := (1 - x_1, \ldots, 1 - x_p)$ for any $\boldsymbol{x} = (x_1, \ldots, x_p) \in \mathbb{C}^p$. This is the main result Corollary 3.7 of this paper. Moreover, we shall determine the differences of $c_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}^{t_1,\ldots,t_{p-1}}(n_1, \ldots, n_r)$ in Corollary 4.3 explicitly.

2 The differences of multiple sequences

In the present section, we study some fundamental properties of the differences of multiple sequences. For this purpose, we use formal power series belonging to $\mathbb{C}[[X]] := \mathbb{C}[[X_1, \ldots, X_r]]$ or $\mathbb{C}[[X, Y]] := \mathbb{C}[[X_1, \ldots, X_r, Y_1, \ldots, Y_r]]$, where r is a positive integer and is fixed throughout this section. We put $\partial_{X_i} = \partial/\partial X_i$ and $\partial_{Y_i} = \partial/\partial Y_i$ for $1 \leq i \leq r$. In the following, we denote the set of non-negative integers by $\mathbb{N}$. We sometimes write n (resp. k) for a sequence $n_1, \ldots, n_r$ (resp. $k_1, \ldots, k_r$) for abbreviation. First, we note that for a formal power series

$$F = \sum_{n,k=0}^{\infty} a(n_1, \ldots, n_r, k_1, \ldots, k_r) \frac{X_1^{n_1} \cdots X_r^{n_r} Y_1^{k_1} \cdots Y_r^{k_r}}{n_1! \cdots n_r!\, k_1! \cdots k_r!} \in \mathbb{C}[[X, Y]],$$

we have the equality

$$(\partial_{X_i} + \partial_{Y_i} - 1)F = \sum_{n,k=0}^{\infty} \Big\{ a(n_1, \ldots, n_i+1, \ldots, n_r, k) + a(n, k_1, \ldots, k_i+1, \ldots, k_r)$$
$$- a(n, k) \Big\} \frac{X_1^{n_1} \cdots X_r^{n_r} Y_1^{k_1} \cdots Y_r^{k_r}}{n_1! \cdots n_r!\, k_1! \cdots k_r!} \quad (3)$$

for any $1 \leq i \leq r$.

Lemma 2.1. *If a formal power series* $F \in \mathbb{C}[[X, Y]]$ *satisfies two conditions*

$$(\partial_{X_i} + \partial_{Y_i} - 1)F = 0 \quad (\textit{for any } 1 \leq i \leq r)$$

and

$$F(X_1, \ldots, X_r, 0, \ldots, 0) = 0,$$

then we have $F = 0$*.*

Proof. If we put

$$F = \sum_{n,k=0}^{\infty} a(n_1, \ldots, n_r, k_1, \ldots, k_r) \frac{X_1^{n_1} \cdots X_r^{n_r} Y_1^{k_1} \cdots Y_r^{k_r}}{n_1! \cdots n_r!\, k_1! \cdots k_r!},$$

by (3) we have

$$a(n_1, \ldots, n_i + 1, \ldots, n_r, k) + a(n, k_1, \ldots, k_i + 1, \ldots, k_r) - a(n, k) = 0$$

for any $n_1, \ldots, n_r, k_1, \ldots, k_r \in \mathbb{N}$ and any $1 \leq i \leq r$. Moreover we have

$$a(n_1, \ldots, n_r, 0, \ldots, 0) = 0$$

for any $n_1, \ldots, n_r \in \mathbb{N}$. By induction on $k_1 + \cdots + k_r$, we see that

$$a(n_1, \ldots, n_r, k_1, \ldots, k_r) = 0$$

holds for any $n_1, \ldots, n_r, k_1, \ldots, k_r \in \mathbb{N}$. □

Definition 2.2. We denote by $\mathbb{C}^{\mathbb{N}^r}$ the set of all mappings from $\mathbb{N}^r$ to $\mathbb{C}$. For each $1 \leq i \leq r$, we define the difference operator $\Delta_i \colon \mathbb{C}^{\mathbb{N}^r} \to \mathbb{C}^{\mathbb{N}^r}$ by putting

$$(\Delta_i a)(n_1, \ldots, n_r) = a(n_1, \ldots, n_r) - a(n_1, \ldots, n_i + 1, \ldots, n_r)$$

for any $a \in \mathbb{C}^{\mathbb{N}^r}$ and any $n_1, \ldots, n_r \in \mathbb{N}$.

We note that $\Delta_i \Delta_j = \Delta_j \Delta_i$ for any $1 \leq i, j \leq r$. We denote the composition

$$\underbrace{\Delta_1 \circ \cdots \circ \Delta_1}_{n_1} \circ \underbrace{\Delta_2 \circ \cdots \circ \Delta_2}_{n_2} \circ \cdots \circ \underbrace{\Delta_r \circ \cdots \circ \Delta_r}_{n_r}$$

by $\Delta_1^{n_1} \Delta_2^{n_2} \cdots \Delta_r^{n_r}$.

Definition 2.3. We define the inversion operator $\nabla \colon \mathbb{C}^{\mathbb{N}^r} \to \mathbb{C}^{\mathbb{N}^r}$ by putting

$$(\nabla a)(n_1, \ldots, n_r) = (\Delta_1^{n_1} \cdots \Delta_r^{n_r} a)(0, \ldots, 0)$$

for any $a \in \mathbb{C}^{\mathbb{N}^r}$ and any $n_1, \ldots, n_r \in \mathbb{N}$.

For a sequence $a\colon \mathbb{N}^r \to \mathbb{C}$, we define

$$f_a = \sum_{n=0}^{\infty} a(n_1, \ldots, n_r) \frac{X_1^{n_1} \cdots X_r^{n_r}}{n_1! \cdots n_r!} \in \mathbb{C}[[X]]$$

and

$$F_a = \sum_{n,k=0}^{\infty} (\Delta_1^{k_1} \cdots \Delta_r^{k_r} a)(n_1, \ldots, n_r) \frac{X_1^{n_1} \cdots X_r^{n_r} Y_1^{k_1} \cdots Y_r^{k_r}}{n_1! \cdots n_r!\, k_1! \cdots k_r!} \in \mathbb{C}[[X, Y]].$$

We note that

$$F_a(X_1, \ldots, X_r, 0, \ldots, 0) = f_a(X_1, \ldots, X_r) \tag{4}$$

and

$$F_a(0, \ldots, 0, Y_1, \ldots, Y_r) = f_{\nabla a}(Y_1, \ldots, Y_r). \tag{5}$$

By (3), we have

$$(\partial_{X_i} + \partial_{Y_i} - 1) F_a = 0 \tag{6}$$

for any $1 \leq i \leq r$.

Proposition 2.4. *Let $a\colon \mathbb{N}^r \to \mathbb{C}$ be a sequence. Then we have*

$$F_{\nabla a}(X_1, \ldots, X_r, Y_1, \ldots, Y_r) = F_a(Y_1, \ldots, Y_r, X_1, \ldots, X_r).$$

Proof. By (6), we have

$$(\partial_{X_i} + \partial_{Y_i} - 1) F_{\nabla a}(X_1, \ldots, X_r, Y_1, \ldots, Y_r) = 0$$

and

$$(\partial_{X_i} + \partial_{Y_i} - 1) F_a(Y_1, \ldots, Y_r, X_1, \ldots, X_r) = 0$$

for any $1 \leq i \leq r$. In addition, by (4) and (5) we have

$$F_{\nabla a}(X_1, \ldots, X_r, 0, \ldots, 0) = f_{\nabla a}(X_1, \ldots, X_r)$$

and

$$F_a(0, \ldots, 0, X_1, \ldots, X_r) = f_{\nabla a}(X_1, \ldots, X_r).$$

Therefore Lemma 2.1 implies the proposition. □

Corollary 2.5. *Let $a\colon \mathbb{N}^r \to \mathbb{C}$ be a sequence. Then for any $n_1, \ldots, n_r, k_1, \ldots, k_r \in \mathbb{N}$, we have*

$$(\Delta_1^{k_1} \cdots \Delta_r^{k_r} (\nabla a))(n_1, \ldots, n_r) = (\Delta_1^{n_1} \cdots \Delta_r^{n_r} a)(k_1, \ldots, k_r).$$

Proof. By comparing the coefficients of both sides of the equation in Proposition 2.4, we obtain the corollary. □

Corollary 2.6. *The operator ∇^2 is the identity on $\mathbb{C}^{\mathbb{N}^r}$.*

Proof. It follows from Corollary 2.5 on setting $n_1 = \cdots = n_r = 0$. □

In the rest of this section, we give explicit expressions for the differences and the inversions of multiple sequences.

Proposition 2.7. *For any sequence* $a\colon \mathbb{N}^r \to \mathbb{C}$*, we have*

$$F_a = f_a(X_1 - Y_1, \ldots, X_r - Y_r)e^{Y_1+\cdots+Y_r}.$$

Proof. It is easily seen that for any $1 \le i \le r$ the right-hand side becomes zero if we apply $\partial_{X_i} + \partial_{Y_i} - 1$. Noting (4) and (6), we get the proposition by Lemma 2.1. $\square$

Corollary 2.8. *For any sequence* $a\colon \mathbb{N}^r \to \mathbb{C}$ *and any* $n_1, \ldots, n_r$*,* $k_1, \ldots, k_r \in \mathbb{N}$*, we have*

$$(\Delta_1^{k_1} \cdots \Delta_r^{k_r} a)(n_1, \ldots, n_r) = \sum_{i_1=0}^{k_1} \cdots \sum_{i_r=0}^{k_r} (-1)^{i_1+\cdots+i_r} \binom{k_1}{i_1} \cdots \binom{k_r}{i_r} a(n_1 + i_1, \ldots, n_r + i_r).$$

Proof. If we apply the operator $\partial_{Y_1}^{k_1} \cdots \partial_{Y_r}^{k_r}$ to the right-hand side of the equation in Proposition 2.7, the result is

$$\sum_{i_1=0}^{k_1} \cdots \sum_{i_r=0}^{k_r} (-1)^{i_1+\cdots+i_r} \binom{k_1}{i_1} \cdots \binom{k_r}{i_r} f_a^{(i_1,\ldots,i_r)}(X_1 - Y_1, \ldots, X_r - Y_r)e^{Y_1+\cdots+Y_r}$$

by the Leibniz rule, where we have put

$$f_a^{(i_1,\ldots,i_r)} = \partial_{X_1}^{i_1} \cdots \partial_{X_r}^{i_r} f_a.$$

Therefore by applying the operator $\partial_{X_1}^{n_1} \cdots \partial_{X_r}^{n_r} \partial_{Y_1}^{k_1} \cdots \partial_{Y_r}^{k_r}$ to both sides of the equation in Proposition 2.7 and comparing the constant term, we obtain the desired equality. $\square$

Corollary 2.9. *Let* $a\colon \mathbb{N}^r \to \mathbb{C}$ *be a sequence. For any* $n_1, \ldots, n_r \in \mathbb{N}$*, we have*

$$(\nabla a)(n_1, \ldots, n_r) = \sum_{i_1=0}^{n_1} \cdots \sum_{i_r=0}^{n_r} (-1)^{i_1+\cdots+i_r} \binom{n_1}{i_1} \cdots \binom{n_r}{i_r} a(i_1, \ldots, i_r).$$

Proof. It follows from Corollary 2.8 on setting $n_1 = \cdots = n_r = 0$. $\square$

3 A proof of the duality

Throughout this section, we fix positive integers r and p. First of all, we define and calculate the inversions of operators on $\mathbb{C}[[X]] := \mathbb{C}[[X_1, \ldots, X_r]]$. Then, we shall prove a generalization of the duality for MHS by using this notion. We begin with the definition of the inversions of formal power series. For a formal power series

$$f = \sum_{n=0}^{\infty} a(n_1, \dots, n_r) \frac{X_1^{n_1} \cdots X_r^{n_r}}{n_1! \cdots n_r!} \in \mathbb{C}[[X]],$$

we define its inversion by

$$\nabla f = \sum_{n=0}^{\infty} (\nabla a)(n_1, \dots, n_r) \frac{X_1^{n_1} \cdots X_r^{n_r}}{n_1! \cdots n_r!}.$$

By Corollary 2.6, the inversion operator ∇ on $\mathbb{C}[[X]]$ is an involution.

Proposition 3.1. *For any formal power series* $f \in \mathbb{C}[[X]]$*, we have*

$$\nabla f = f(-X_1, \dots, -X_r) e^{X_1 + \cdots + X_r}.$$

Proof. It is immediate from (5) and Proposition 2.7. □

For a mapping $\xi \colon \mathbb{C}[[X]] \to \mathbb{C}[[X]]$, we define its inversion by

$$(\nabla \xi) f = \nabla(\xi(\nabla f)), \quad f \in \mathbb{C}[[X]].$$

Clearly, we have

$$\nabla(\xi f) = (\nabla \xi)(\nabla f)$$

for any $\xi \colon \mathbb{C}[[X]] \to \mathbb{C}[[X]]$ and any $f \in \mathbb{C}[[X]]$ since the inversion operator ∇ on $\mathbb{C}[[X]]$ is an involution. It is easily seen that the equality

$$\nabla(\xi \eta) = (\nabla \xi)(\nabla \eta) \tag{7}$$

holds for any mappings ξ, $\eta \colon \mathbb{C}[[X]] \to \mathbb{C}[[X]]$. We note that the inversion operator ∇ is a linear operator.

Proposition 3.2. *For any* $1 \le i \le r$*, we have*

$$\nabla X_i = -X_i \quad \textit{and} \quad \nabla \partial_{X_i} = 1 - \partial_{X_i},$$

where X_i *is the operator on* $\mathbb{C}[[X]]$ *defined by* $f \mapsto X_i f$ *for any* $f \in \mathbb{C}[[X]]$*.*

Proof. By Proposition 3.1, for any $f \in \mathbb{C}[[X]]$ we have

$$(\nabla X_i)(\nabla f) = \nabla(X_i f) = -X_i f(-X_1, \dots, -X_r) e^{X_1 + \cdots + X_r} = -X_i(\nabla f),$$

which implies $\nabla X_i = -X_i$. If we calculate $(\nabla \partial_{X_i}) f$ by definition, the other assertion also follows easily. □

In the following, we consider a generalization of the duality for MHS. We recall that we have fixed positive integers r and p. Let $\boldsymbol{x}_i = (x_{i1}, \dots, x_{ip}) \in$

$\mathbb{C}^p$ for $1 \le i \le r$ and let $t_1, \ldots, t_{p-1} \in \mathbb{C} \setminus \{0, -1, -2, \ldots\}$. Then for any $n_1, \ldots, n_r \in \mathbb{N}$, we define $c_{\boldsymbol{x}_1; \cdots ; \boldsymbol{x}_r}^{t_1, \ldots, t_{p-1}}(n_1, \ldots, n_r)$ by

$$\sum_{\substack{n_1 = n_{11} \ge \cdots \ge n_{1p} \ge 0 \\ \cdots \\ n_r = n_{r1} \ge \cdots \ge n_{rp} \ge 0}} \frac{\displaystyle\prod_{i=1}^{r} \binom{n_i}{\nu_{i1}, \ldots, \nu_{ip}} x_{i1}^{\nu_{i1}} \cdots x_{ip}^{\nu_{ip}}}{\displaystyle\prod_{j=1}^{p-1} \binom{n_{1j} + \cdots + n_{rj} + t_j - 1}{\nu_{1j} + \cdots + \nu_{rj}} (n_{1j} + \cdots + n_{rj} + t_j)},$$

where we have put

$$\nu_{i1} = n_{i1} - n_{i2}, \quad \ldots, \quad \nu_{ip-1} = n_{ip-1} - n_{ip}, \quad \nu_{ip} = n_{ip}$$

for abbreviation. In the case $p = 1$, we have

$$c_{x_1; \cdots ; x_r}(n_1, \ldots, n_r) = x_1^{n_1} \cdots x_r^{n_r}. \tag{8}$$

The following Proposition 3.3 is the key to proving the duality for the nested sums $c_{\boldsymbol{x}_1; \cdots ; \boldsymbol{x}_r}^{t_1, \ldots, t_{p-1}}(n_1, \ldots, n_r)$. For any $\boldsymbol{x} = (x_1, \ldots, x_p) \in \mathbb{C}^p$ with $p \ge 2$, we define

$$^{-}\boldsymbol{x} = (x_2, \ldots, x_p) \in \mathbb{C}^{p-1}.$$

Proposition 3.3. *Let* $\boldsymbol{x}_i = (x_{i1}, \ldots, x_{ip}) \in \mathbb{C}^p$ *for* $1 \le i \le r$ *and let* $t_1, \ldots, t_{p-1} \in \mathbb{C} \setminus \{0, -1, -2, \ldots\}$. *If* $p \ge 2$, *for any* $n_1, \ldots, n_r \in \mathbb{N}$ *we have*

$$\begin{aligned} &(n_1 + \cdots + n_r + t_1) c_{\boldsymbol{x}_1; \cdots ; \boldsymbol{x}_r}^{t_1, \ldots, t_{p-1}}(n_1, \ldots, n_r) \\ &\quad - \sum_{k=1}^{r} x_{k1} n_k c_{\boldsymbol{x}_1; \cdots ; \boldsymbol{x}_r}^{t_1, \ldots, t_{p-1}}(n_1, \ldots, n_k - 1, \ldots, n_r) = c_{^{-}\boldsymbol{x}_1; \cdots ; ^{-}\boldsymbol{x}_r}^{t_2, \ldots, t_{p-1}}(n_1, \ldots, n_r). \end{aligned} \tag{9}$$

Proof. Let $1 \le k \le r$. In the proof we write $\sum_\sharp$ and $\sum_\flat$ instead of

$$\sum_{\substack{n_1 \ge n_{12} \ge \cdots \ge n_{1p} \ge 0 \\ \cdots \\ n_k > n_{k2} \ge \cdots \ge n_{kp} \ge 0 \\ \cdots \\ n_r \ge n_{r2} \ge \cdots \ge n_{rp} \ge 0}} \quad \text{and} \quad \sum_{\substack{n_1 \ge n_{12} \ge \cdots \ge n_{1p} \ge 0 \\ \cdots \\ n_r \ge n_{r2} \ge \cdots \ge n_{rp} \ge 0 \\ \exists i, \, n_i > n_{i2}}},$$

respectively. The kth term in the second sum of the left-hand side of (9) is equal to

$$\sum_\sharp R_k \times \frac{x_{k1}^{\nu_{k1}} \cdots x_{kp}^{\nu_{kp}} \displaystyle\prod_{\substack{1 \le i \le r \\ i \ne k}} \binom{n_i}{\nu_{i1}, \ldots, \nu_{ip}} x_{i1}^{\nu_{i1}} \cdots x_{ip}^{\nu_{ip}}}{\displaystyle\prod_{j=2}^{p-1} \binom{n_{1j} + \cdots + n_{rj} + t_j - 1}{\nu_{1j} + \cdots + \nu_{rj}} (n_{1j} + \cdots + n_{rj} + t_j)}, \tag{10}$$

where

$$R_k = n_k \binom{n_k - 1}{\nu_{k1} - 1, \nu_{k2}, \ldots, \nu_{kp}} \binom{n_1 + \cdots + n_r + t_1 - 2}{\nu_{11} + \cdots + \nu_{r1} - 1}^{-1} \frac{1}{n_1 + \cdots + n_r + t_1 - 1}.$$

By a simple calculation we obtain

$$R_k = \frac{\nu_{k1}}{\nu_{11} + \cdots + \nu_{r1}} \binom{n_k}{\nu_{k1}, \ldots, \nu_{kp}} \binom{n_1 + \cdots + n_r + t_1 - 1}{\nu_{11} + \cdots + \nu_{r1}}^{-1},$$

which is obviously equal to 0 if $n_k = n_{k2}$ (i.e. $\nu_{k1} = 0$). Therefore we can replace $\sharp$ by $\flat$ in the expression (10). Consequently the second sum of the left-hand side of (9) is equal to

$$\sum_{\flat} \frac{\displaystyle\prod_{i=1}^{r} \binom{n_i}{\nu_{i1}, \ldots, \nu_{ip}} x_{i1}^{\nu_{i1}} \cdots x_{ip}^{\nu_{ip}}}{\displaystyle\binom{n_1 + \cdots + n_r + t_1 - 1}{\nu_{11} + \cdots + \nu_{r1}} \prod_{j=2}^{p-1} \binom{n_{1j} + \cdots + n_{rj} + t_j - 1}{\nu_{1j} + \cdots + \nu_{rj}} (n_{1j} + \cdots + n_{rj} + t_j)},$$

from which the desired equality is easily derived. □

In order to rewrite Proposition 3.3 in terms of generating functions, we introduce the following operator on $\mathbb{C}[[X]]$

$$\xi_{x_1,\ldots,x_r} = X_1 \partial_{X_1} + \cdots + X_r \partial_{X_r} - x_1 X_1 - \cdots - x_r X_r$$

for complex numbers $x_1, \ldots, x_r$. For this operator we have

$$\nabla \xi_{x_1,\ldots,x_r} = \xi_{1-x_1,\ldots,1-x_r} \tag{11}$$

by Proposition 3.2. We note that for a formal power series

$$f = \sum_{n=0}^{\infty} a(n_1, \ldots, n_r) \frac{X_1^{n_1} \cdots X_r^{n_r}}{n_1! \cdots n_r!} \in \mathbb{C}[[X]]$$

and a complex number t, the equality

$$\begin{aligned} (\xi_{x_1,\ldots,x_r} + t) f = \sum_{n=0}^{\infty} \Big\{ & (n_1 + \cdots + n_r + t) a(n_1, \ldots, n_r) \\ & - \sum_{k=1}^{r} x_k n_k a(n_1, \ldots, n_k - 1, \ldots, n_r) \Big\} \frac{X_1^{n_1} \cdots X_r^{n_r}}{n_1! \cdots n_r!} \end{aligned} \tag{12}$$

holds.

Lemma 3.4. *For any* $x_1, \ldots, x_r \in \mathbb{C}$ *and any* $t \in \mathbb{C} \setminus \{0, -1, -2, \ldots\}$, *the operator* $\xi_{x_1,\ldots,x_r} + t$ *on* $\mathbb{C}[[X]]$ *is an injection.*

Proof. Let

$$f = \sum_{n=0}^{\infty} a(n_1, \ldots, n_r) \frac{X_1^{n_1} \cdots X_r^{n_r}}{n_1! \cdots n_r!} \in \mathbb{C}[[X]]$$

be a formal power series in the kernel of the operator $\xi_{x_1,\dots,x_r}+t$. Then, by (12), the equality

$$(n_1+\cdots+n_r+t)a(n_1,\dots,n_r)=\sum_{k=1}^{r}x_kn_ka(n_1,\dots,n_k-1,\dots,n_r)$$

holds for any $n_1,\dots,n_r\in\mathbb{N}$. Since we have

$$n_1+\cdots+n_r+t\neq 0$$

for any $n_1,\dots,n_r\in\mathbb{N}$, we obtain $f=0$ by induction on $n_1+\cdots+n_r$. ☐

For any $\boldsymbol{x}_1,\dots,\boldsymbol{x}_r\in\mathbb{C}^p$ and any $t_1,\dots,t_{p-1}\in\mathbb{C}\setminus\{0,-1,-2,\dots\}$, we put

$$f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}=\sum_{n=0}^{\infty}c^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}(n_1,\dots,n_r)\frac{X_1^{n_1}\cdots X_r^{n_r}}{n_1!\cdots n_r!}\in\mathbb{C}[[X]].$$

In the case $p=1$, we have

$$f_{x_1;\cdots;x_r}=e^{x_1X_1+\cdots+x_rX_r} \tag{13}$$

by (8).

Proposition 3.5. *Let* $\boldsymbol{x}_i=(x_{i1},\dots,x_{ip})\in\mathbb{C}^p$ *for* $1\le i\le r$ *and let* $t_1,\dots,t_{p-1}\in\mathbb{C}\setminus\{0,-1,-2,\dots\}$. *If* $p\ge 2$ *then we have*

$$(\xi_{x_{11},\dots,x_{r1}}+t_1)f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}=f^{t_2,\dots,t_{p-1}}_{^-\boldsymbol{x}_1;\cdots;^-\boldsymbol{x}_r}.$$

Proof. It follows from Proposition 3.3 and (12). ☐

Now we state the main result of this paper. We put

$$\boldsymbol{1}-\boldsymbol{x}=(1-x_1,\dots,1-x_p)$$

for any $\boldsymbol{x}=(x_1,\dots,x_p)\in\mathbb{C}^p$.

Theorem 3.6. *For any* $\boldsymbol{x}_1,\dots,\boldsymbol{x}_r\in\mathbb{C}^p$ *and any* $t_1,\dots,t_{p-1}\in\mathbb{C}\setminus\{0,-1,-2,\dots\}$, *we have*

$$\nabla f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}=f^{t_1,\dots,t_{p-1}}_{\boldsymbol{1}-\boldsymbol{x}_1;\cdots;\boldsymbol{1}-\boldsymbol{x}_r}.$$

Proof. By Proposition 3.5 and (13), we have

$$(\xi_{x_{1p-1},\dots,x_{rp-1}}+t_{p-1})\cdots(\xi_{x_{11},\dots,x_{r1}}+t_1)f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}=e^{x_{1p}X_1+\cdots+x_{rp}X_r}.$$

Applying the inversion operator ∇ to both sides, we obtain

$$\begin{aligned}(\xi_{1-x_{1p-1},\dots,1-x_{rp-1}}+t_{p-1})\cdots(\xi_{1-x_{11},\dots,1-x_{r1}}+t_1)(\nabla f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r})&\\ =e^{(1-x_{1p})X_1+\cdots+(1-x_{rp})X_r}&\end{aligned}$$

by (7), (11) and Proposition 3.1. On the other hand we have

$$\begin{aligned}(\xi_{1-x_{1p-1},\dots,1-x_{rp-1}}+t_{p-1})\cdots(\xi_{1-x_{11},\dots,1-x_{r1}}+t_1)f^{t_1,\dots,t_{p-1}}_{\boldsymbol{1}-\boldsymbol{x}_1;\cdots;\boldsymbol{1}-\boldsymbol{x}_r}&\\ =e^{(1-x_{1p})X_1+\cdots+(1-x_{rp})X_r},&\end{aligned}$$

which together with Lemma 3.4 implies the theorem. ☐

Corollary 3.7. *For any* $\boldsymbol{x}_1,\dots,\boldsymbol{x}_r\in\mathbb{C}^p$ *and any* $t_1,\dots,t_{p-1}\in\mathbb{C}\backslash\{0,-1,-2,\dots\}$, *we have*

$$\nabla c^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}=c^{t_1,\dots,t_{p-1}}_{\boldsymbol{1}-\boldsymbol{x}_1;\cdots;\boldsymbol{1}-\boldsymbol{x}_r}.$$

Proof. It is clear from Theorem 3.6. □

4 The difference formula

In this section, we shall prove a formula which gives explicitly the differences of the nested sums $c^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}(n_1,\dots,n_r)$. As usual, we put

$$[\xi,\eta]=\xi\eta-\eta\xi$$

for mappings ξ, $\eta\colon\mathbb{C}[[X]]\to\mathbb{C}[[X]]$.

Lemma 4.1. *Let* $i_1,\dots,i_q$ *be distinct integers satisfying* $1\le i_1,\dots,i_q\le r$ *and let* $x_1,\dots,x_r,t$ *be complex numbers. If we put*

$$x_{i_1}+\cdots+x_{i_q}=c,$$

then we have

$$[\partial_{X_{i_1}}+\cdots+\partial_{X_{i_q}}-c,\,\xi_{x_1,\dots,x_r}+t]=\partial_{X_{i_1}}+\cdots+\partial_{X_{i_q}}-c.$$

Proof. The left-hand side is equal to

$$[\partial_{X_{i_1}},\,X_{i_1}\partial_{X_{i_1}}]+\cdots+[\partial_{X_{i_q}},\,X_{i_q}\partial_{X_{i_q}}]-x_{i_1}[\partial_{X_{i_1}},\,X_{i_1}]-\cdots-x_{i_q}[\partial_{X_{i_q}},\,X_{i_q}].$$

Since we have $[\partial_{X_j},\,X_j]=1$ and $[\partial_{X_j},\,X_j\partial_{X_j}]=\partial_{X_j}$ for all $1\le j\le r$, we obtain the assertion. □

Theorem 4.2. *Let* $\boldsymbol{x}_1,\dots,\boldsymbol{x}_r\in\mathbb{C}^p$ *and let* $c\in\mathbb{C}$. *We suppose that*

$$\boldsymbol{x}_{i_1}+\cdots+\boldsymbol{x}_{i_q}=\underbrace{(c,\dots,c)}_{p}$$

for distinct integers $1\le i_1,\dots,i_q\le r$. *Then for any* $t_1,\dots,t_{p-1}\in\mathbb{C}\setminus\{0,-1,-2,\dots\}$, *we have*

$$(\partial_{X_{i_1}}+\cdots+\partial_{X_{i_q}}-c)f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}=0.$$

Proof. The proof is by induction on p. The case $p=1$ follows directly from (13). Let $p\ge 2$. We put $\boldsymbol{x}_i=(x_{i1},\dots,x_{ip})$ for any $1\le i\le r$. Since we have

$$(\xi_{x_{11},\dots,x_{r1}}+t_1+1)(\partial_{X_{i_1}}+\cdots+\partial_{X_{i_q}}-c)=(\partial_{X_{i_1}}+\cdots+\partial_{X_{i_q}}-c)(\xi_{x_{11},\dots,x_{r1}}+t_1)$$

by Lemma 4.1, it holds that

$$(\xi_{x_{11},\dots,x_{r1}}+t_1+1)(\partial_{X_{i_1}}+\cdots+\partial_{X_{i_q}}-c)f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}=0$$

by Proposition 3.5 and the hypothesis of induction. This, together with Lemma 3.4, leads to

$$(\partial_{X_{i_1}}+\cdots+\partial_{X_{i_q}}-c)f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}=0.$$

Therefore we complete the proof. □

Corollary 4.3. *Let* $\boldsymbol{x}_1, \dots, \boldsymbol{x}_r \in \mathbb{C}^p$ *and let* $t_1, \dots, t_{p-1} \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$*. Then, for any* $n_1, \dots, n_r, k_1, \dots, k_r \in \mathbb{N}$*, we have*

$$(\Delta_1^{k_1} \cdots \Delta_r^{k_r} c^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r})(n_1, \dots, n_r) = c^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r;\mathbf{1}-\boldsymbol{x}_1;\cdots;\mathbf{1}-\boldsymbol{x}_r}(n_1, \dots, n_r, k_1, \dots, k_r).$$

Proof. By Theorem 4.2, we have

$$(\partial_{X_i} + \partial_{Y_i} - 1) f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r;\mathbf{1}-\boldsymbol{x}_1;\cdots;\mathbf{1}-\boldsymbol{x}_r}(X_1, \dots, X_r, Y_1, \dots, Y_r) = 0$$

for any $1 \le i \le r$. We also have

$$f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r;\mathbf{1}-\boldsymbol{x}_1;\cdots;\mathbf{1}-\boldsymbol{x}_r}(X_1, \dots, X_r, 0, \dots, 0) = f^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}(X_1, \dots, X_r),$$

which results from

$$c^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r;\mathbf{1}-\boldsymbol{x}_1;\cdots;\mathbf{1}-\boldsymbol{x}_r}(n_1, \dots, n_r, 0, \dots, 0) = c^{t_1,\dots,t_{p-1}}_{\boldsymbol{x}_1;\cdots;\boldsymbol{x}_r}(n_1, \dots, n_r).$$

Therefore by Lemma 2.1, (4) and (6) we obtain the assertion. ☐

By this corollary, we see that the differences of the nested sums considered in this paper are again the same kind of nested sums. Corollary 3.7 follows also from Corollary 4.3 on setting $n_1 = \cdots = n_r = 0$.

References

[1] D. M. Bradley, Duality for finite multiple harmonic q-series, Disc. Math. **300** (2005), 44-56.

[2] M. Hoffman, Quasi-symmetric functions and mod p multiple harmonic sums, preprint arXiv:math.NT/0401319.

[3] G. Kawashima, A class of relations among multiple zeta values, J. Number Theory. **129** (2009), 755-788.

[4] G. Kawashima and T. Tanaka, Newton series and extended derivation relations for multiple L-values, preprint arXiv:0801.3062.

[5] J. A. M. Vermaseren, Harmonic sums, Mellin transforms and integrals, Int. J. Mod. Phys. A **14** (1999), 2037-2076.

[6] J. Zhao, Wolstenholme type theorem for multiple harmonic sums, Int. J. Number Theory **4** (2008), 73-106.

Graduate School of Mathematics
Nagoya University
Chikusa-ku, Nagoya 464-8602, Japan
E-mail: m02009c@math.nagoya-u.ac.jp